

SISTER CHROMATID EXCHANGES: A STUDY IN FLUOROTIC INDIVIDUALS OF NORTH GUJARAT

F J Sheth, A S Multani and N J Chinoy
Ahmedabad, India

SUMMARY: The purpose of this preliminary investigation was to compare the genotoxic effect of fluoride in human individuals directly exposed to high concentrations of drinking water fluoride (1.95 to 2.2 ppm) with those in individuals exposed to concentrations (0.6 to 1.0 ppm) within the WHO permissible limit. Sister chromatid exchanges (SCE) and cell cycle proliferative index were studied in whole-blood cultures of fluorotic individuals and compared with normal controls of age-matched groups. The results suggest that the SCE rate in persons exposed to fluoride in the endemic (1.95-2.2 ppm) areas of North Gujarat is significantly higher than that of those living in Ahmedabad (0.6-1.0 ppm). There is no significant difference in the cell cycle proliferative index. Further investigations are needed to confirm the findings.

Key words: Fluorosis; Genotoxic; Mutagen; Sister chromatid exchanges.

Introduction

Fluoride is ubiquitously distributed in our environment. It is present in air, water, and fertilizer residues, and traces have been reported in almost every edible material that we consume in our daily life. Water is ordinarily the principal source of fluoride intake by the human population. Continuous exposure to higher than permissible levels of fluoride (> 1.0 ppm) in drinking water leads to crippling fluorosis. More than a million people in India are afflicted with skeletal and dental fluorosis. Numerous studies have been carried out in relation to the influence of fluoride on the environmental and biological systems, but there is a great deal of controversy regarding its potential genotoxic effects in mammals (1,2). The present study was undertaken with a view to investigate the genotoxic effect of fluoride, if any, on the endemic population of Gujarat.

Materials and Methods

Peripheral blood was collected from 100 individuals, mainly from Mehsana District of North Gujarat, and 21 from Ahmedabad. The inhabitants of these villages were examined for apparent mottled teeth, back pain, stiffness of back and joints, and other clinical manifestations including skeletal problems prior to blood sampling. Most of these individuals were unable to bend due to a stiff back, which indicated that they were afflicted with fluorosis. The clinical history was recorded in a standard proforma. Water samples of the villages were collected for the estimation of fluoride levels.

Metaphase chromosome preparations from the blood samples of the fluoride-affected individuals and control members were carried out by routine phytohaemagglutinin (PHA) stimulated cultures. 5-Bromodeoxyuridine (BrdU) was added at a final concentration of 10 $\mu\text{g/ml}$ at 0 hours.

Division of Human Genetics, Department of Zoology, University School of Sciences, Gujarat University, Ahmedabad 380 009, India. Correspondence to Professor Chinoy.

The fluorescence plus Giemsa method was used for the scoring of sister chromatid exchanges, and the procedure of Perry and Wolff (3) was followed with slight modifications. All the slides were randomized with code numbers and were scanned by a single individual. The chromosomes were examined under a Nikon light microscope. The influence on the cell cycle proliferative index was evaluated by recording the percentage of cells in the first (M1), second (M2), and third (M3) cell divisions. A minimum of 100 cells were analysed from each culture. The cells in M2 phase were further examined and only the cells with well spread chromosomes were scored for sister chromatid exchanges (SCEs). A total of 25 cells from each culture were examined for SCE frequency. The Students 't' test was used for statistical analysis of SCE data.

Fluoride concentrations from the serum as well as from the water samples were determined with an Ion Selective Electrode Orion Model 701A.

Results

The frequencies of sister chromatid exchanges in fluoride affected individuals and in normal controls are shown in Table 1. The values are the average (mean) number of SCEs observed in minimum 25 M2 cells each with 46 chromosomes. Statistical analysis indicated a significant increase ($p < 0.05$) in the SCE/plate in the study group when compared to the normal controls. The influence of fluoride toxicity on cell cycle proliferative index was also carried out. Cultures from both the fluoride affected individuals and normal controls showed presence of cells in 1st, 2nd and 3rd cell cycles indicating the existence of all three cell cycles in both the groups of cells. No significant variation was observed in the cell cycle proliferative index of fluoride affected individuals as compared to controls (Figure 1).

The serum fluoride level was significantly elevated ($p < 0.001$) in persons residing in Gujarat as compared to those examined in Ahmedabad (Table 2).

Discussion

A number of investigators have utilized the SCE test to study the genotoxicity of fluoride (1,4-12). In the present study, human populations directly exposed to fluoride in drinking water in endemic regions of North Gujarat were investigated to evaluate the possible effect of fluoride on SCE. To the best of our knowledge this is the first report on genotoxic effects following long-term fluoride intake in an endemic area in India. In this study, the results from cultured human lymphocytes indicate a significant increase in the frequency of SCE in the study group compared to the normal control. *In vitro* studies have yielded inconsistent findings with regard to SCE in white blood cells. In human lymphocytes exposed to doses up to 4 mM (11) and 3 mM (12) of NaF, no significant variation in the frequency of SCE was reported. *In vivo* studies carried out in mice maintained for seven generations on a diet containing 1.2 mM NaF indicated no increase in SCE frequencies in bone marrow cells (8). On the other hand, an increase in the incidence of SCE was observed in cultured Red Muntjac cells and in cultured Syrian Hamster embryo cells at dosage levels of 0.5, 1.0, and 1.9 mM, respectively (9,10).

TABLE 1. Frequencies of SCEs in control and study group

Sr.No.	Group	No. of cases investigated	No. of metaphases screened	No. of metaphases with 2nd cell cycle	SCE/plate*
1	Control	21	2173	612	7.46 ± 0.52
2	Study	100	11005	3856	8.79 ± 0.26

* = mean ± SE

The values are significantly different ($p < 0.05$).

TABLE 2. Fluoride concentration in drinking water samples and in serum of control and fluorotic cases screened for SCE


Area	Water Fluoride (ppm)	Serum Fluoride (ppm)
Control (Ahmedabad)	0.62 ± 0.02	0.04 ± 0.002
Range	0.60 - 1.04	0.03 - 0.05
n	24	25
Endemic Region (Mehsana District)	2.20 ± 0.05*	0.286 ± 0.02*
Range	1.949 - 2.211	0.177 - 0.539
n	36	24

Values are mean ± SE

n = number of samples

* = $p < 0.001$

FIGURE: Influence of fluoride on the cell cycle proliferation



Furthermore, no significant chromosomal damage was observed in lymphocytes of cattle with signs of chronic fluoride poisoning (13), in mouse oocytes at doses as high as 100 ppm, and ewe oocytes exposed to 10 ppm NaF (14). A significant increase in chromosome aberration was reported in human leukocytes (15), human fibroblasts (16,17), and in mice (18) maintained on fluoridated water.

It is, however, difficult to correlate the results of the present study with the earlier reports, since most of them have been carried out using either *in vitro* cultures or *in vivo* systems by treating the experimental animal with fluoridated compounds. Moreover, the doses of fluoride used for *in vitro* studies were much higher than those ever attained *in vivo* in body fluids of mammalian organisms (19), and the *in vivo* studies have been carried out at concentrations far in excess of those encountered in daily living (2). In the present study, the fluoride content in drinking water of the endemic villages (Table 2) was higher than the permissible level of 1 ppm, according to the WHO standards (20). Serum fluoride concentrations were also significantly higher in the fluorotic subjects of these areas, corroborating earlier data (21-23).

The results of the present investigation suggest that in fluoride-affected persons exposed to 1.95-2.2 ppm fluoride in drinking water chromosomal alterations as indicated by SCE frequency and chromosome aberrations were higher than in normal persons exposed to 0.6-1.0 ppm drinking water fluoride. However, analysis of samples from other endemic areas, and also use of a series of investigations including cytogenetic and mutagenic assays, are necessary to confirm the findings. Work is also needed on a large control group who are consuming 0 ppm or less than 0.2 ppm fluoride. However, the fluoride levels in drinking water in Ahmedabad and its vicinity range from 0.6 to 1.04 ppm (see Table 2).

Acknowledgements

This work was supported by the Council of Scientific and Industrial Research, New Delhi. The help rendered by Shri Nanak Kumar, Engineer, Sewage and Water Board, Sidhpur, and Dr Mahendrabhai A Patel in collection of blood samples during the field visits is gratefully acknowledged.

References

- 1 Li Y, Heerema NA, Dunipace AJ, Stookey GK. Genotoxic effects of fluoride evaluated by sister chromatid exchange. *Mutation Research* 192 191-201 1987.
- 2 Li Y, Dunipace AJ, Stookey GK. Genotoxic effects of fluoride: a controversial issue. *Mutation Research* 195 127-136 1988.
- 3 Perry P, Wolff S. New Giemsa method for the differential staining of sister chromatids. *Nature* 251 156-158 1974.
- 4 Wolff S. Sister chromatid exchange. *Annual Review of Genetics* 11 183-201 1977.
- 5 Abe S, Motomichi S. SCE as an index of mutagenesis and/or carcinogenesis. In: Sandberg A (Ed). *Sister Chromatid Exchange*. Liss, New York 1982 pp 461-514.
- 6 Perry P, Thomson EJ. The methodology of sister chromatid exchanges. In: Kilbey BJ, Legator M, Nichols W, Ramel C (Eds). *Handbook of Metagenicity Test Procedures*. 2nd ed. Elsevier, Amsterdam 1984 pp 495-529.
- 7 Perry P and Evans HJ. Cytological detection of mutagen-carcinogen exposure by sister chromatid exchange. *Nature* 258 121-125 1975.

- 8 Kram D, Schneider EL, Singer L, Martin GR. The effects of high and low fluoride diets on the frequencies of sister chromatid exchanges. *Mutation Research* 57 51-55 1978.
- 9 He W, Liu H, Bao H *et al.* Effect of sodium fluoride and fluoroacetamide on sister chromatid exchanges and chromosomal aberrations in cultured Red Muntjac (*Muntjac muntjak*) cells. *Acta Scientiae Circumstantiae* 3 94-100, 1983.
- 10 Tsutsui T, Susuki N, Ohmori M. Sodium fluoride-induced morphological and neoplastic transformation. Chromosome aberrations, sister chromatid exchanges, and unscheduled DNA synthesis in cultured Syrian hamster embryo cells. *Cancer Research* 44 938-941 1984.
- 11 Kishi K, Tonomura A. Cytogenetic effects of sodium fluoride. *Mutation Research* 130 367 1984.
- 12 Thomson EJ, Kilanowski FM, Perry PE. The effect of fluoride on chromosome aberration and sister chromatid exchange frequencies in cultured human lymphocytes. *Mutation Research* 144 89-92 1985.
- 13 Leonard A, Duknuds GH, Decat G, Leonard ED. Cytogenetic investigations on leukocytes of cattle intoxicated with fluoride. *Toxicology* 7 239-242 1977.
- 14 Jagiello G, Lin JS. Sodium fluoride as potential mutagen in mammalian eggs. *Archives of Environmental Health* 29 230-235 1974.
- 15 Jachimczak D, Skotarczak B. The effect of fluoride and lead ions on the chromosomes of human leucocytes in vitro. *Genetica Polonica* 19 353-357 1978.
- 16 Tsutsui T, Suzuki N, Ohmori M, Maizumi H. Cytotoxicity, chromosome aberrations and unscheduled DNA synthesis in cultured human diploid fibroblasts induced by sodium fluoride. *Mutation Research* 139 193-198, 1984.
- 17 Scott D. Cytogenetic effects of sodium fluoride in cultured human fibroblasts. 4th International Conference on Environmental Mutagens, Stockholm, 1985.
- 18 Mohamed AH, Chandler, ME. Cytological effects of sodium fluoride on mice. *Fluoride* 15 (3) 110-118 1982.
- 19 Katz S, McDonald Jr JL, Stookey GK. *Preventive Dentistry in Action*. DCP, Upper Montclair NJ 1979 pp 306-392.
- 20 *Fluorine and Fluorides. Environmental Health Criteria* 36. World Health Organization, Geneva 1984 pp 1-136.
- 21 Chinoy NJ, Narayana MV, Sequeira E *et al.* Studies on effects of fluoride in 36 villages of Mehsana District, North Gujarat. *Fluoride* 25 101-110 1992.
- 22 Chinoy NJ, Narayana MV. Studies on fluorosis in Mehsana District of North Gujarat. *Proceedings, Zoological Society, Calcutta* 45 157-161 1992.
- 23 Chinoy NJ, Barot VV, Michael M *et al.* Fluoride toxicity studies in Mehsana District, North Gujarat. *Journal of Environmental Biology* 15 163-173 1994.