Fluoride Vol. 31 No. 4 203-216 1998. Research Report 203

# AMELIORATION OF FLUORIDE TOXICITY BY VITAMINS E AND D IN REPRODUCTIVE FUNCTIONS OF MALE MICE

N J Chinoy\* and Arti Sharma Ahmedabad, India

SUMMARY: Studies on the beneficial effects of vitamins E and D supplementation on functions of caput and cauda epididymides, their spermatozoa, vas deferens and seminal vesicle of sodium fluoride (NaF) treated (10 mg/kg body weight) male mice (Mus musculus) were carried out. The NaF treatment resulted in significant decrease in the body and epididymis weight but those of vas deferens and seminal vesicle were not affected. NaF treatment brought about alterations in epididymal milieu as elucidated by the significant decrease in levels of sialic acid and protein as well as activity of ATPase in epididymides. As a result, the sperm maturation process was affected leading to a significant decline in cauda epididymal sperm motility and viability. This caused a significant reduction in fertility rate. The cauda epididymal sperm count was also significantly reduced. The data obtained suggest that fluoride treatment induced significant metabolic alterations in the epididymides, vas deferens and seminal vesicles of mice. The withdrawal of NaF treatment (30 days) produced incomplete recovery. On the other hand, sup-plementation of vitamins E or D during the withdrawal period of NaF treated mice was found to be very beneficial in recovery of all NaF induced effects, thus elucidating their ameliorative role in recovery from toxic effects of NaF on the reproductive functions and fertility. On the whole, a combination of vitamins E and D treatment was comparatively more effective than that with vitamin E or D alone. Therefore, vitamin therapy could be beneficial for the amelioration of fluoride induced changes in reproductive functions.

Key words. Epididymis; Fluoride toxicity; Reversibility; Seminal vesicle; Vas deferens; Vitamin D; Vitamin E.

### INTRODUCTION

Fluoride is one of the potent toxicants to which humans are exposed. Extensive data on skeletal and dental fluorosis are available.<sup>1</sup> However, the effect of fluoride on the structure and metabolism of several soft tissues has been reported recently. Messer *et al*<sup>2</sup> reported that low levels of fluoride in food rendered mice infertile, while a high fluoride diet improved their fertility. These reports were contradicted by Tao and Suttie,<sup>3</sup> whose experiments showed that fluoride did not play any essential role in reproduction. Later Kour and Singh<sup>4</sup> reported that the testicular spermatogenic process was affected in mice administered fluoride at a dose of 500 and 1000 ppm in drinking water. Li *et al*<sup>5</sup> claimed that fluoride did not have adverse effects on spermatogenesis or sperm morphology. Earlier, our laboratory reported that ingestion of 10 or 20 mg sodium fluoride in mice caused alterations in the histology of reproductive organs, morphology of sperm and induced biochemical changes.<sup>6,7</sup>

Recent investigations<sup>8,9</sup> showed that fluoride interferes with the structural and functional integrity of testis, internal milieu of epididymis, vas deferens and also affected the metabolism and morphology of spermatozoa of mice, rats and rabbits and reduces fertility. The above results clearly indicate that some

<sup>\*</sup>Address for correspondence. Reproductive Endocrinology and Toxicology Unit, UGC Department of Special Assistance and COSIST in Zoology, School of Sciences, Gujarat University, Ahmedabad 380009, India.

reproductive organs. The present study was undertaken because of the above conflicting reports and paucity of data on the toxic effects of fluoride on male reproductive system and its possible reversal by vitamins E and D.

### MATERIALS AND METHODS

Healthy, adult male mice (*Mus musculus*) of Swiss strain, weighing between 25 to 30 g obtained from the National Institute of Occupational Health, Ahmedabad, were used for the experiments. They were housed in an air conditioned animal house at a temperature of  $26\pm2^{\circ}$ C and exposed to 10 to 12 hours of daylight.

The animals were divided into seven different groups (see protocol in Table 1) and caged separately. Group I (control) mice were maintained on standard diet and water ad libitum. Group II (control) mice were fed olive oil orally with vitamins E and D dissolved in the oil. Sodium fluoride (NaF) (Loba Chemie, Bombay, 99% purity) was administered orally to Group III-VII mice at a dose of 10 mg/kg body weight for 30 days using a feeding tube attached to a hypodermic syringe. The dose of NaF was based on the LD<sub>50</sub> of male mice which is 54.4 mg  $F^-$  /kg body weight.<sup>10</sup> Group III mice were sacrificed on the 31<sup>st</sup> day. NaF treatment of Groups IV-VII animals was withdrawn after 30 days and the animals were maintained on standard diet and water ad libitum for a further 30 days to study any reversibility of the induced effects. Additionally, during this 30 day withdrawal period, Group V mice were administered vitamin E (Roche Products Ltd, Bombay), Group VI mice received vitamin D<sub>3</sub> (Teva Pharmaceuticals Ind. Ltd, Israel), and Group VII animals received a combination of vitamins D and E. The doses of vitamins D and E were 0.002  $\mu$ g/day/animal and 2  $\mu$ g/day/animal respectively. These doses were based on earlier studies.

	Table 1. Experimenta	111010001		
Group	Treatment and dose	Duration (days)	Day of autopsy	No. of animals used
I	Untreated control	-	Sacrificed with treated	20
11	Vehicle (olive oil) treated control	-	Sacrificed with treated	20
111	Sodium fluoride (NaF, 10 mg/kg body weight)	30	31 <sup>st</sup>	20
IV	NaF as in Group III, on 31st day NaF withdrawal for further 30 days	30 + 30	61 <sup>st</sup>	20
V	NaF as in Group III, on 31st day NaF withdrawal + vit.E for further 30 days	30 + 30	61 <sup>st</sup>	20
VI	NaF as in Group III, on 31st day NaF withdrawal + vit.D for further 30 days	30 + 30	61 <sup>st</sup>	20
VII	NaF as in Group III, on 31st day NaF withdrawal + vit.E + vit.D for further 30 days	30 + 30	61 <sup>st</sup>	20

The control and treated animals were sacrificed by cervical dislocation. The caput and cauda epididymides, vas deferens and seminal vesicle were carefully dissected out, blotted free of blood, weighed on a Roller Smith (USA) torsion balance and utilized for the various parameters as follows:

Amelioration of F<sup>-</sup> toxicity by vitamins D and E 205

# **Biochemical studies**

Sperm motility and count: The cauda epididymal sperm suspension was prepared in normal saline. The percent motility and count of cauda epididymal spermatozoa of normal and all treated groups of mice were determined by the method of Prasad *et al*<sup>11</sup> and expressed as percentage motility and millions/ml respectively.

Sperm viability: The ratio of live:dead spermatozoa of control and all treated groups of mice were determined using 1% trypan blue as described in the method of Talbot and Chacon.<sup>12</sup> The number of stained and unstained spermatozoa were scored in 10-20 separate fields and percentages (%) were calculated.

*Fertility test:* Conducted by cohabiting estrous or proestrous female animals with treated male animals in a ratio of 2:1 on the 31st day of treatment according to the WHO protocol MB50.<sup>13</sup> The vaginal smear was checked the next morning for the presence of sperm, which indicated that mating had occurred. This was day one of pregnancy. The animals were autopsied on day 16 of pregnancy and the implantation sites were counted and compared with the control.

Protein: The protein levels of caput and cauda epididymides and vas deferens of control and all treated animals were estimated by the method of Lowry et  $al^{14}$  at 540 nm on a Spectronic-88 Bausch and Lomb spectrophotometer. Protein was expressed as mg/100 mg fresh tissue weight.

Adenosine Triphosphatase (ATPase) (E.C. 3.6.1.3): The ATPase activity in epididymides of control and all treated groups of mice were assayed following the method of Quinn and White<sup>15</sup>. The activity was expressed as  $\mu$  moles of ip released/mg protein/30 minutes.

Sialic acid: Sialic acid in the epididymis of both control and treated animals was determined by the method of Jourdian *et al*<sup>16</sup> and was expressed as  $\mu$ g/mg fresh tissue weight.

Phosphorylase (E.C. 2.4.1.1): Phosphorylase activity in vas deferens of control and treated mice was determined by the method of Cori *et al*<sup>17</sup> and the enzyme activity was expressed as  $\mu g$  phosphorus released/mg protein/15 minutes.

Glycogen: The glycogen concentrations in vas deferens of control and treated mice were estimated by the method of Seifter *et al*<sup>18</sup> and expressed as  $\mu g$  glycogen/100 mg fresh tissue weight.

*Fructose:* Fructose concentration in the seminal vesicles was estimated by the method of Foreman *et al*<sup>19</sup> and was expressed as  $\mu$ g/mg tissue weight.

*Statistics:* For all biochemical estimations, a minimum of 8 to 10 replicates were done for each parameter. The data were statistically analysed using student's t test and Analysis of Variance (ANOVA) followed by Scheffe's test for least significance.

## RESULTS

Body weight: The body weight of the NaF treated Group III animals decreased (P<0.001) after 30 days of treatment in comparison to both the control Groups I-II. After 30 days of NaF withdrawal, the body weight of Group IV mice did not recover as compared to control Group I. On administration of vitamins E or D alone and in combination, the body weights of Groups V-VII animals recovered significantly (P<0.001) compared to control Group II (Table 2).

Organ weight: The weights of caput and cauda epididymides in Group III mice declined significantly (P<0.01) compared to control Groups I-II. In Group IV (withdrawal) the recovery was less and the difference was not significant as compared to control, whereas significant recovery was obtained in Group V (vit E), VI (vit D) and VII (vit E + vit D). However, weights of seminal vesicle and vas deferens were not affected by different treatments (Table 2).

			Organ Weights (mg)						
Groups	Body Weight (g)	Caput Epididymis	Cauda Epididymis	Vas deferens	Seminal Vesicle				
1	35.6 ± 0.19	14.0 ± 0.17	12.0 ± 0.4	11.6 ± 0.35	56.0 ± 0.37				
H	36.5 ± 0.21	14.8 ± 0.33	12.6 ± 0.6	12.0 ± 0.20	56.9 ± 0.29				
HI	30.4 ± 0.24**	11.2 ± 0.30*	9.0 ± 0.5*	13.2 ± 0.33 <sup>NS</sup>	58.4 ± 0.16 <sup>NS</sup>				
IV	33.5 ± 0.20 <sup>NS</sup>	13.0 ± 0.28 <sup>NS</sup>	10.6 ± 0.4 <sup>NS</sup>	12.4 ± 0.21 <sup>NS</sup>	57.2 ± 0.21 <sup>NS</sup>				
V	37.6 ± 0.23**	15.0 ± 0.28*	13.8 ± 0.17*	12.8 ± 0.23 <sup>NS</sup>	57.0 ± 0.19 <sup>NS</sup>				
VI	37.2 ± 0.20**	14.2 ± 0.20 <sup>NS</sup>	13.4 ± 0.21*	12.6 ± 0.19 <sup>NS</sup>	57.6 ± 0.22 <sup>NS</sup>				
VII	37.8 ± 0.26**	15.2 ± 0.23*	13.9 ± 0.13*	11.9 ± 0.16*	57.9 ± 0.17 <sup>NS</sup>				

TABLE 2.	Body (g) an	d organ weight	(mg) of control	and treated	groups of mice
----------	-------------	----------------	-----------------	-------------	----------------

Values are Mean  $\pm$  S.E. \*P<0.01 \*\*P<0.001 NS = Non Significant Groups: I = Untreated Control; II = Vehicle (Olive oil) treated control; III = NaF treated; IV = NaF withdrawal for 30 days; V = NaF as in III, withdrawal on day 31 and vit E for further 30 days; VI = NaF as in III, withdrawal on day 31 and vit D for further 30 days; VII = NaF as in III, withdrawal on day 31 and vits E + D feeding for further 30 days

Sperm motility: The motility of cauda epididymal sperm of Group III NaF treated mice decreased significantly (P<0.001) as compared to control Groups I-II. At the end of the withdrawal phase, sperm motility recovered in Groups IV-VII, but was still lower than control Groups I-II. Group IV had significant but relatively low recovery (P<0.01). Sperm motility of Group V on vitamin E, and Group VII on vitamins E and D recovered comparatively better than Group VI on vitamin D, but all three groups recovered significantly better (P<0.001) than Group IV. (Tables 3A and 3B).

Sperm count: The sperm count in the cauda epididymis of Group III NaF treated mice declined significantly (P<0.001) compared to control Groups I-II. In Group IV (withdrawal) significant (P<0.01) recovery was obtained, whereas vitamin E or D treated animals showed very significant (P<0.001) recovery as compared to the NaF treated (Group III) mice which was again more with vitamin E. In Group VII the recovery was most significant (P<0.001) (Tables 3A and 3B).

Sperm viability: Cauda epididymal sperm viability (live:dead ratio) was significantly reduced (P<0.001) by NaF treatment as compared to control groups I-II. However, significant (P<0.001) recovery was obtained in Groups IV, V and VI (withdrawal, vit E, vit D). In the latter two groups, recovery was more than in Group IV. In Group VII almost complete recovery was noted and was comparable to normal values (Table 3A).

Parameters	Group I Intreated Control	Group II Vehicle treated	Group III NaF	Group IV Withdrawał	Group V Vitamin E	Group VI Vitamin D	Group VII Vit E + Vit D
Sperm motility (%)	75.28 ± 1.58	76.19 ± 1.34	26.31 ± 0.55**	43.73 ± 0.79*	67.42 ± 0.73**	52.46 ± 0.91**	68.66 ± 0.92**
Sperm count (millions/ml)	42.33 ± 0.45	43 ± 0.81	22 ± 0.81**	28.6 ± 0.55*	35.5 ± 1.70**	31.66 ± 1.37**	38.0 ± 0.59**
Sperm viability ( Live : dead	%) 72.15:27.85 ± 0.50	73.62:26.38 ± 0.54	13.89:86.11 ± 0.44**	39.67:60.33 ± 0.31**	66.52:33.48 ± 0.42**	59.23:40.77 ± 0.33**	69.98:30.0 ± 0.39**
Fertility Rate (%)	95 - 100 +ve	95 - 100 +ve	0**	27.97 ± 3.4** +ve	72.31 ± 0.77** +ve	65.47 ± 2.36** +ve	78.13 ± 0.93** +ve
Values are M	lean + S E * 1	2<0.01 ** P<0.0	$01 \rightarrow ve = necitive$				

Table 3A.	Cauda epidid	lymal sperm motility,	count, viability	y and fertilit	y rate of control	and treated	groups of mice
-----------	--------------	-----------------------	------------------	----------------	-------------------	-------------	----------------

Values are Mean ± S.E. \* P<0.01 \*\* P<0.001 +ve = positive Groups I and II compared to Group III (NaF) Group III (NaF) compared to Groups IV, V, VI, VII

TABLE 3B. ANO	VA of cauda	epididymal	sperm motil	ity and count
---------------	-------------	------------	-------------	---------------

Source of variation	df	SS	MSS	f(cal)	f(tab)
Sperm motility	· · · · · · · · · · · · · · · · · · ·				
Groups	6	12629.82	2104.97	40.40	0.00
Residual	35	5495.46	157.01	13.40	2.30
Sperm count					
Group	6	2048.41	341.40	070.02	0.06
Residual	35	42.74	1.22	219.83	2.20

Significance at 5% level

df = Degree of Freedom SS = Sum of Squares MSS = Mean Sum of Square

Fertility rate: Fluoride treatment led to a significant (P<0.001) inhibition of fertility rate as compared to control Groups I-II. The fertility was restored significantly (P<0.001) upon withdrawal of treatment for 30 days as compared to NaF treated animals (Group III), but ingestion of vitamin E or D in Groups V and VI resulted in very significant (P<0.001) restoration of fertility which was more in Group V (vit E) and in Group VII (vit E + vit D) (Table 3A).

**Protein:** The protein levels in the caput and cauda epididymides and vas deferens decreased significantly (P<0.001) after 30 days of NaF treatment in Group III as compared to control Groups I-II. The protein levels in Group IV (withdrawal) showed insignificant recovery after 30 days except in cauda epididymis where recovery was comparatively more (P<0.01). However, administration of vitamins E or D (Groups V-VI) resulted in significant recovery in protein levels in caput and cauda epididymides (P<0.001) and vas deferens (P<0.01). In Group VII the protein levels recovered to almost the same levels as in control mice (Tables 4A, 4B).

ATPase: Activity of ATPase in caput and cauda epididymides of NaF treated mice showed a significant (P<0.001) decrease in comparison to both the control Groups I-II. The recovery was less in Group IV (withdrawal) but significant (P<0.001) recovery was obtained in caput and cauda epididymides in Groups V and VI (vitamins E and D treated). Combination of vitamin E + vitamin D treatment brought about more recovery than vitamin E or vitamin D alone (Tables 5A and 5B).

Sialic acid: Levels of sialic acid in caput and cauda epididymides declined significantly (P<0.01) after NaF treatment as compared to both the control Groups I and II. However, by withdrawal of treatment (Group IV) and subsequent administration of vitamin E or D (Groups V, VI), significant (P<0.01) recovery was noted as compared to control mice. The recovery was most significant (P<0.001) in Group VII where combination of vit E and vit D was given (Tables 6A and 6B).

Glycogen: A significant (P<0. 001) accumulation of glycogen in the vas deferens of NaF treated mice was observed as compared to control Groups I-II. In Group IV the recovery was not significant. On the other hand, significant recovery was obtained after vitamin E ingestion (Group V) (P<0.001) and in Group VI (P<0.01) as compared to NaF treated mice. However, in Group VII (vit E and vit D) significant (P<0.001) recovery was observed (Tables 7A and 7B).

Phosphorylase: NaF treatment resulted in significant (P<0.001) suppression of vas deferens phosphorylase activity as compared to control Groups I-II. However, after withdrawal of treatment and administration of vitamins E or D (Groups V-VI), the activity of phosphorylase was restored back to normal (Groups IV, V, VI). The recovery was comparatively less in Group IV but was significant (P<0.001) with both vitamin treatments. Combined treatment of vit E and vit D was the most effective in restoring enzyme activity to almost normal levels (Tables 7A and 7B).

*Fructose:* Fructose levels in seminal vesicle increased significantly (P<0.001) after NaF treatment for 30 days as compared to control Groups I-II. The recovery was significant (P<0.01) by withdrawal (Group IV) of treatment. On administration of vitamins E or D (Groups V-VI) the levels of fructose recovered significantly (P<0.001). Moreover, the levels of fructose in vit E + vit D treated mice (Group VII) were also significantly (P<0.001) restored to normal as compared to NaF treated mice (Tables 7A and 7B).

Organs	Group I Untreated Control	Group II Vehicle treated	Group III NaF	Group IV Withdrawal	Group V Vitamin E	Group VI Vitamin D	Group VII Vit E + Vit D
Caput Epididymis	11.43 ± 0.28	12.55 ± 0.14	8.47 ± 0.20**	9.25 ± 0.16 <sup>NS</sup>	11.00 ± 0.24**	10.61 ± 0.19**	12.0 ± 0.18**
Cauda Epididymis	14.10 ± 0.19	15.18 ± 0.37	8.52 ± 0.44**	9.89 ± 0.12*	12.92 ± 0.33**	11.48 ± 0.16**	13.62 ± 0.29**
Vas Deferens	14.09 ± 0.09	14.43 ± 0.21	8.87 ± 0.20**	9.82 ± 0.17 <sup>NS</sup>	11.79 ± 0.26*	11.77 ± 0.15*	12.98 ± 0.18**

# **TABLE 4A.** Protein levels (mg/100 mg fresh tissue weight) in caput and cauda epididymides and vas deferens of control and treated groups of mice

Values are Mean ± S.E. \* P<0.01 \*\* P<0.001 NS = Non Significant

Groups I and II compared to Group III (NaF)

Group III (NaF) compared to Groups IV, V, VI, VII

		•	• •			
Source of variation	df	SS	MSS	f(cal)	f(tab)	
Caput epididymal protein						
Groups	6	81.75	13.62	29.04	2.26	
Residual	35	10.86	0.35	30.91	2.30	
Cauda epididymal proteir	 ו		· · · · · · · · · · · · · · · · · · ·		<u> </u>	
Groups	6	2047.43	341.23	404 53	2.26	
Residual	35	24.49	0.69	494.00	2.30	
Vas deferens protein		<u></u>			<u></u>	
Group	6	135.94	22.65	066 47	244	
Residual	28	2.38	0.085	200.47	2.44	

TABLE 4B.	ANOVA of c	aput and	cauda e	pididymis	and vas	s deferens	protein

Significance at 5% level

df = Degrees of Freedom SS = Sum of Squares MSS = Mean Sum of Square

. .

in caput and cauda epididymides of control and treated groups of mice											
Organs	Group I Untreated Control	Group II Vehicle treated	Group III NaF	Group IV Withdrawal	Group V Vitamin E	Group VI Vitamin D	Group VII Vit E + Vit D				
Caput Epididymis	1.99 ± 0.04	2.01 ± 0.06	0.93 ± 0.07**	1.10 ± 0.03 <sup>NS</sup>	1.69 ± 0.08*	1.65 ± 0.06*	1.78 ± 0.05**				
Cauda Epididymis	2.08 ± 0.07	2.12 ± 0.09	0.978 ± 0.03**	1.23 ± 0.05*	1.87 ± 0.08**	1.79 ± 0.10**	1.96 ± 0.08**				

.....

Values are Mean ± S.E. \* P<0.01 \*\* P<0.001 NS = Non Significant Groups I and II compared to Group III (NaF)

Group III (NaF) compared to Groups IV, V, VI, VII

Source of variation	df	SS	MSS	f(cal)	f(tab)
Caput epididymal ATPas	e				
Groups	6	5.42	0.90	10.00	0.40
Residual	28	1.53	0.05	10.00	2.42
Cauda epididymal ATPa	se			······································	· · · · · · · · · · · · · · · · · · ·
Group	6	5.82	0.97	4.64.68	2.40
Residual	28	0.17	0.006	00.101	2.42
			······		

TABLE 5B. ANOVA of caput and cauda epididymal ATPase

Significance at 5% level

df = Degrees of Freedom

SS = Sum of Squares MSS = Mean Sum of Square

TABLE 6	6A.	Sialic acid	levels	(µg/m	g tissue	wt) i	n caput and	cauda	epididymides	of contro	I and treate	d groups of	mice
---------	-----	-------------	--------	-------	----------	-------	-------------	-------	--------------	-----------	--------------	-------------	------

Organs	Group I Untreated Control	Group II Vehicle treated	Group III NaF	Group IV Withdrawal	Group V Vitamin E	Group VI Vitamin D	Group VII Vit E + Vit D
Caput Epididymis	4.72 ± 0.03	4.89 ± 0.02	3.76 ± 0.07*	4.14 ± 0.02 <sup>NS</sup>	4.45 ± 0.05*	4.34 ± 0.02*	4.60 ± 0.06**
Cauda Epididymis	5.56 ± 0.06	5.59 ± 0.09	4.48 ± 0.07*	5.05 ± 0.04*	5.37 ± 0.04*	5.18 ± 0.02*	5.47 ± 0.03**

Values are Mean ± S.E. \* P< 0.01 \*\* P< 0.001 NS = Non Significant Groups I and II compared to Group III (NaF) Group III (NaF) compared to Groups IV, V, VI, VII

TABLE 6B. ANOVA of cauda and caput epididymal sialic acid

Source of variation	df	SS	MSS	f(cal)	f(tab)
Caput epididymis					
Groups	6	5.59	0.93	24.05	2.26
Residual	35	1.11	0.03	31.05	2.30
Cauda epididymis					
Groups	6	6.32	1.05	07 77	2.26
Residual	35	0.45	0.012	01.11	2.30

Significance at 5% level

df = Degrees of Freedom SS = Sum of Squares MSS = Mean Sum of Square

Organe	Parameters	Group	Group II	Group III	Group IV	Group V	Group VI	Group VII
organs		Untreated Control	Vehicle treated	NaF	Withdrawal	Vitamin E	Vitamin D	Vit E + Vit D
Vas Deferens	Glycogen (µg/100mg fresh tissue wt)	776.54 ± 20.65	815.85 ± 19.76	1281.29 ± 15.36* *	1181.29 ± 5.36 <sup>NS</sup>	858.35 ± 11.56* *	1058.86 ± 19.92*	829.67 ± 13.63**
	Phosphorylase (µg phosphorus/mg protein/15 min)	53.17 ± 1.45	54.0 ± 1.32	36.76 ± 0.88**	42.30 ± 1.01*	51.07 ± I.75**	45.22 ± 1.06**	52.01 ± 1.68**
Seminal Vesicle	Fructose (µg/mg tissue wt)	14.84 ± i.11	14.98 ± 1.16	20.83 ± 1.20**	17.62 ± 0.99*	16.37± 0.90**	16.97 ± 1.0l**	15.23 ± 0.93**

# TABLE 7A Showing alycogen concentration and phosphorylase activity in vas deferens

Values are Mean ± S.E. NS = Non significant \* P< 0.01 \*\* P< 0.001

Source of variation	df	SS	MSS	f(cal)	f(tab)
Glycogen (vas deferens)					
Groups	6	1293659.4	215609.9	8 20	2.44
Residual	28	735336.6	26262.02	0.20	
Phosphorylase (vas defe	erens)				
Groups	6	5697.17	949.52	5.06	2.42
Residual	28	4458.4	159.22	5.90	
Fructose (seminal vesicl	e)				
Group	6	123.12	20.52	19 /9	2.42
Residual	28	31.19	1.11	10.40	

#### TABLE 78 ANOVA of glycogen phosphorylase and fructose

Significance at 5% level df = Degrees of Freedom SS = Sum of Squares MSS = Mean Sum of Square

### Amelioration of F<sup>-</sup> toxicity by vitamins D and E 213

#### DISCUSSION

The present investigation was carried out to explore the effects of fluoride (NaF) and the possible ameliorative role of vitamins E and D ingestion on epididymis, vas deferens and seminal vesicle of mice during withdrawal period.

The sodium fluoride treatment caused a decrease in the body weight. Similar results were reported by others in rats and mice fed with different concentrations of fluoride.<sup>8,20</sup> An insignificant reduction in weight of epididymis occurred but those of seminal vesicle and vas deferens were not affected.

In the present study the levels of protein in caput and cauda epididymides and vas deferens showed a significant decrease after 30 days of NaF treatment. This decrease might be due to impairment of protein metabolism/synthesis. Earlier studies carried out from our laboratory and elsewhere have also reported a dose dependent decrease in protein levels in the serum and reproductive organ of mice, rats and rabbits.<sup>7,8,21</sup>

Sialic acid is an important constituent of mucopolysaccharides and sialomucoproteins which are essential for the maturation of spermatozoa in epididymis and maintenance of the structural integrity of their membranes.<sup>22</sup> The levels of sialic acid in caput and cauda epididymides were decreased in the present study. Hence it is likely that the structural integrity of acrosomal membrane of the sperm might have been altered. A decline in sialic acid concentration of NaF treated mouse and rat epididymis has also been reported earlier.<sup>8</sup>

Adenosine triphosphatase activity in caput and cauda epididymides showed a decline with greater propensity in cauda epididymis. Alterations in the activity of ATPase in NaF treated mouse and rat epididymis and rabbit spermatozoa have been reported.<sup>8,21</sup> It was reported<sup>23</sup> that fluoride acts directly on the motile apparatus without substantially affecting other metabolic pathways as it inhibits the dynein ATPase in sperm. According to Hodge and Smith,<sup>24</sup> NaF toxicity involves inhibition of enzyme activity, particularly those in which divalent metal cations act as co-factors. In the present study too, the alterations in ATPase activity might be related to the fact that it is either a Mg<sup>2+</sup> or a Ca<sup>2+</sup> activated enzyme.

Reports from our laboratory have revealed that the glycogen concentrations were enhanced in vas deferens of fluoride treated rats and mice.<sup>7,25</sup> These results are in agreement with the observations of the present investigation, wherein glycogen was found to accumulate in vas deferens of NaF treated mice. The increase in glycogen could be correlated with the decrease in the activity of phosphorylase in the vas deferens by NaF ingestion. Fluoride has been reported to alter carbohydrate metabolism mainly by causing allosteric inhibition of some key enzymes in glycolysis and tricarboxylic acid (TCA) cycle.<sup>26</sup>

The increase in level of fructose in seminal vesicle after 30 days of NaF exposure further supports our observation on alteration in carbohydrate metabolism of fluorotic mice. As fructose has a vital role in providing energy to the sperm, it is evident that the increased fructose level might influence sperm metabolism.

Fluoride is known to inhibit sperm motility, glycolysis and respiration process. It has been demonstrated<sup>27</sup> that bovine sperm treated with 30 mM fluoride became immobile within two minutes. Human spermatozoa lost their motility *in vitro* in the presence of 250 mM NaF within 20 minute incubation.<sup>28</sup> In the present study too, a significantly low sperm motility was obtained by NaF ingestion.

The sperm density of cauda epididymis in NaF treated group of mice also declined significantly. The decrease could be correlated with the testicular spermatogenic arrest following fluoride ingestion in mice, rats and rabbits.<sup>8,21</sup>

The evaluation of NaF treated spermatozoa stained with trypan blue showed a large number of dead sperm, probably due to loss of membrane permeability, which might be another major factor in the decline in sperm motility.<sup>25</sup>

The above mentioned alterations in sperm motility, density and metabolism might be the outcome of the altered and hostile internal milieu of the epididymis of NaF treated mice since it is known that normal epididymal structure and its internal microenvironment are important for sperm maturation and for maintaining them in a viable, motile state.<sup>22,29,30</sup>

The reduction in sperm motility, count, viability and changes in their metabolism led to the significant decline in fertility of treated mice as also reported earlier in mice, rats and rabbits.<sup>8,21,31</sup>

In the present study, NaF treatment affected numerous androgen dependent parameters in the epididymis, vas deferens and seminal vesicle to a variable extent. Hence, it is evident that fluoride affects male reproductive organs and fertility.

In withdrawal group of animals (Group IV) the NaF induced effects were not restored completely to normal state after 30 days. However, in Groups V, VI, VII of animals treated with vitamin E or vitamin D alone and in combination in the withdrawal period, almost complete recovery from fluoride toxicity was obtained. The extent of recovery was more pronounced with vitamin E as compared to D and was most significant with the combined treatment.

In rats, the main symptoms of vitamin E deficiency are degeneration of the testis, abnormalities of gestation, regression in the ovary and changes in ovulation.<sup>32</sup> At the cellular-molecular level, vitamin E is believed to exert its protective effect primarily through destruction of cell-damaging free-radical oxygen species.<sup>33</sup> In the present study too, vitamin E ingestion was beneficial for recovery from fluoride toxicity.

According to earlier reports, elevated levels of vitamin D mitigated the symptoms of fluorosis. Studies carried out by Gupta *et al* 1996<sup>34</sup> have revealed that the treatment of vitamins C, D and calcium showed a significant improvement in the skeletal, clinical fluorosis and biochemical parameters in children consuming water containing 4.5 ppm of fluoride.

The above reports and earlier work carried out by Chinoy and associates have elucidated that therapeutic agents like amino acids, vitamin C and  $Ca^{2+}$  could mitigate fluoride induced effects.<sup>8,35,36</sup>

Thus, in conclusion, sodium fluoride has a definite effect on reproduction. However, the fluoride induced effects are reversible and transient and could be effectively reversed by withdrawal of treatment and subsequent supplementation of vitamins E and D. Thus vitamins E or D may be used as therapeutic agents for the mitigation of fluoride induced toxicity in endemic areas all over the world. Hence, these results have important implications for amelioration of fluorosis in endemic regions.

### Amelioration of F<sup>-</sup> toxicity by vitamins D and E 215

### REFERENCES

- 1 Susheela AK, Koacher J, Jain SK, Sharma K, Jha M. Fluoride toxicity: A molecular approach. In: Susheela AK (Ed). Fluoride toxicity. Proceedings of the 13<sup>th</sup> Conference of the International Society for Fluoride Research. Kalpana Printing House, New Delhi 1985 pp 78-79.
- 2 Messer HH, Armstrong WD, Singer L. Fertility impairment in mice on a low fluoride intake, *Science 177* 893-894 1972.
- 3 Tao S, Suttie JW. Evidence for a lack of an effect of dietary fluoride level on reproduction in mice. *Journal of Nutrition 106* 1115-1122 1976.
- 4 Kour K, Singh J. Histological findings in mice testes following fluoride ingestion. Fluoride 13 (4) 160-162 1980.
- 5 Li Y, Dunipace AT, Stookey GK. Effects of fluoride on the mouse sperm morphology test. *Journal of Dental Research 66* (11) 1509-1511 1987.
- 6 Chinoy NJ, Sequeira E. Effects of fluoride on the histoarchitecture of reproductive organs of the male mouse. *Reproductive Toxicology 3* (4) 261-267 1989a.
- 7 Chinoy NJ, Sequeira E. Fluoride induced biochemical changes in reproductive organs of male mice. *Fluoride 2* (2) 78-85 1989b.
- 8 Chinoy NJ. Role of fluoride in animal systems: A review. In: Toxicity and Monitoring of Xenobiotics. 1995 pp 13-30.
- 9 Chinoy NJ, Sequeira E, Narayana MV. Effects of vitamin C and calcium on the reversibility of fluoride induced alterations in spermatozoa of rabbit. *Fluoride 24* (1) 29-39 1991a.
- Pillai KS, Mathai AT, Deshmukh PB. Acute toxicity of fluoride to mice. *Fluoride 20* (2) 65-70 1987.
- 11 Prasad MRN, Chinoy NJ, Kadam KM. Changes in succinate dehydrogenase level in rat epididymis under normal and altered physiologic conditions. *Fertility and Sterility* 23 180-190 1972.
- 12 Talbot P, Chacon RS. A triple stain technique for evaluating normal acrosome reaction of human sperm. *Journal of Experimental Zoology 215* 201-208 1981.
- 13 WHO Protocol MB-50: A method for examining the effect of a plant extract administered orally on the fertility of male rats. AP/PD, 991 4E 1983.
- 14 Lowry OH, Rosebrough MJ, Farr AL, Randall RJ. Protein measurement with the Folin-Phenol reagent. *Journal of Biological Chemistry* 193 256-275 1951.
- 15 Quinn PJ, White IG. Distribution of adenosine triphosphatase activity in ram and bull spermatozoa. *Journal of Reproduction and Fertility* 15 449-452 1968.
- 16 Jourdian GW, Dean L, Roseman S. A periodate resorcinol method for the quantitative estimation of free sialic acid and their glycosidase. *Journal of Biological Chemistry* 246 (2) 430-435 1971.
- 17 Cori CF Cori GT, Green A. Crystalline muscle phosphorylase kinetics. Journal of Biological Chemistry 151 39-55 1943.
- 18 Seifter S, Dayton S, Novic B, Muntwyler E. The estimation of glycogen with anthrone reagent. Archives of Biochemistry 25 191-200 1950.
- 19 Foreman D, Gaylor L, Evans R, Trella C. A modification of Roe procedure for the determination of fructose with increased specificity. *Analytical Chemistry* 56 584-590 1973.
- 20 Saralakumari D, Varadacharyulu NCh, Ramakrishna Rao P. Effects of fluoride toxicity on growth and lipids in liver, kidney and serum in rats. *Arogya Journal of Health Sciences* 15 24-29 1988.

- 21 Chinoy NJ. Effects of fluoride on physiology of some animals and human beings. Indian Journal of Environment and Toxicology 1 (1) 17-32 1991a.
- 22 Rajalakshmi M, Prasad MRN. Contribution of the epididymis and vas deferens in maturation of spermatozoa. In: Talwar GP (Ed). *Recent Advances in Reproduction and Regulation of Fertility.* Elsevier, Amsterdam 1979 pp 253-258.
- 23 Blum JJ, Hayes A. Specific anion effects on ATPase activity, calmodulin sensitivity and solubilization of dynein ATPase. *Journal of Cellular Biochemistry* 25 197-212 1984.
- 24 Hodge HC, Smith FA. Fluoride. In: Lee DHK (Ed). Metallic Contaminants and Human Health. Academic Press, New York 1972 p 163.
- 25 Chinoy NJ, Narayana MV, Dalal V, Rawat M, Patel D. Amelioration of fluoride toxicity in some accessory reproductive glands and spermatozoa of rat. *Fluoride* 28 (2) 75-86 1995.
- 26 Dousset JC, Rioufol C, Philibert C, Bourbon P. Effects of inhaled HF on cholesterol, carbohydrate and trioxycarboxylic acid metabolism in guinea pigs. *Fluoride* 20 137-141 1987.
- 27 Schoff PI, Lardy MA. Effects of fluoride and caffeine on metabolism of ejaculated bovine spermatozoa. *Biology of Reproduction 37* 1037-1046 1987.
- 28 Chinoy NJ, Narayana MV. In vitro fluoride toxicity in human spermatozoa. Reproductive Toxicology 8 (2) 155-159 1994.
- 29 Chinoy NJ. Structure and function of epididymis in relation to vulnerable points of intervention for male fertility regulation. *Indian Review of Life Sciences* 4 37-68 1984.
- 30 Cooper TG. The Epididymal Sperm Maturation and Fertilization. Springer-Verlag, New York 1986 p 281.
- 31 Chinoy NJ and Sequeira E. Reversible fluoride induced fertility impairment in male mice. *Fluoride 25* (2) 71-76 1992.
- 32 Marks J. A Guide to the Vitamins. Their Role in Health and Disease. University Park Press, Baltimore, Maryland 1975 pp 51-66.
- 33 McCay PD, King MM. Vitamin E: Its role as biological free radical scavenger and its relationship to the microsomal mixed function oxidase system. In: Machlin LJ (Ed). Vitamin E: A comprehensive treatise. Marcel Dekker Inc, New York and Basel 1980 pp 289-317.
- 34 Gupta SK, Gupta RC, Seth AK, Gupta A. Reversal of fluorosis in children. Acta Paediatrica Japonica 385 513-519 1996.
- 35 Chinoy NJ. Effects of fluoride on some organs of rats and their reversal. Proceedings of the Zoological Society (Calcutta) 44 (1) 11-15 1991b.
- 36 Chinoy NJ, Patel D. Ameliorative role of amino acids on fluoride induced alterations in uterine carbohydrate metabolism in mice. *Fluoride 29* (4) 217-226 1996.

PUBLISHED BY THE INTERNATIONAL SOCIETY FOR FLUORIDE RESEARCH Editor's Office: 1620 Massachusetts Street, Lawrence, Kansas 66044, USA