HISTOLOGICAL CHANGES IN THE BRAIN OF YOUNG FLUORIDE-INTOXICATED RATS

YM Shivarajashankara,^a AR Shivashankara,^a P Gopalakrishna Bhat,^b S Muddanna Rao,^c S Hanumanth Rao^d Karnataka, India

SUMMARY: Wistar albino rats were exposed to 30 or 100 ppm fluoride (as NaF) in drinking water during their fetal, weanling, and post-weaning stages until the age of ten weeks. Rats exposed to 30 ppm fluoride did not show any notable alterations in brain histology, whereas rats exposed to 100 ppm fluoride showed significant neurodegenerative changes in the hippocampus, amygdala, motor cortex, and cerebellum. Changes included decrease in size and number of neurons in all the regions, decrease in the number of Purkinje cells in the cerebellum, and signs of chromatolysis and gliosis in the motor cortex. These histological changes suggest a toxic effect of high-fluoride intake during the early developing stages of life on the growth, differentiation, and subcellular organization of brain cells in rats.

Keywords: Albino rats, Amygdala, Brain histology, Cerebellum, Fluoride intoxication, Hippocampus, Motor cortex, Neurodegenerative changes, Sodium fluoride.

INTRODUCTION

Long-term intake of high levels of fluoride in humans causes neurological complications such as paralysis of limbs, vertigo, spasticity in extremities, and impaired mental acuity.¹ Fluoride is also known to cross the blood-brain barrier.² Fluoride accumulation was observed in the brain of experimental animals exposed to chronic high-fluoride intake, and this accumulation increased as the drinking water fluoride levels increased.³ Chronic fluoride toxicity is also known to cause altered neuronal and cerebrovascular integrity,⁴ abnormal behaviour patterns,³ and metabolic lesions in the brain,⁵⁻⁷ in experimental animals.

In the present study, effects of chronic fluoride intoxication on brain cell structure were assessed by exposing rats to 30 or 100 ppm fluoride in drinking water at the early stages of life and then studying the histology of various brain regions.

MATERIALS AND METHODS

Adult (4-6 month-old) wistar albino rats were used in the study. The rats were fed with a standard pelleted diet (Hindustan Lever Ltd., India), and were given water *ad libitum*. The animals were maintained under proper temperature, ventilation, and hygienic conditions.

Pregnant rats were divided into three groups: control, 30F, and 100F. Control rats received drinking water with 0.5 ppm fluoride, while the 30F and 100F rats received 30 tively, in their drinking wa throughout the lactation per rats on weaning and were (0.5 ppm, 30 ppm, and 10 weeks. The ten-week-old rate = 9) groups were sacrificed transcardially with 0.9% so were removed.

The brains were fixed in of the hippocampus, amyg The sections were process stained sections were obser

RE

No histological changes when compared to the corre-100F group, however, sho various brain regions; the l bellum showed neurodegen

In the hippocampus, neusmall nucleus, and there was terations were seen in cornudentate gyrus sub regions of and CA₃ sub regions. The changes among the various long-term intake of high levreported fluoride accumular the hippocampus. The patter findings of Varner *et al*⁴ wh CA₄ areas of the hippocamposed to 1-ppm fluoride in the

Neurodegenerative chang bellum were similar to those motor cortex, neurons in the there were also signs of ch observed chromatin clumpi lation, presence of ghost-like rats exposed to chronic into work, the Purkinje cells in the lation, and there was an imcrease in the number of Purk

^aFor Correspondence: YM Shivarajashankara, Dept. of Biochemistry, MR Medical College, Gulbarga-585105, Karnataka, India; E-mail: shivrajsym@yahoo.com. ^bDept. of Biochemistry, Kasturba Medical College, Manipal-576119, Karnataka, India. ^cDept. of Anatomy, Kasturba Medical College, Manipal-576119, Karnataka, India. ^dDept. of Biochemistry, KBN Institute of Medical Sciences, Gulbarga-585104, Karnataka, India.

and 100F rats received 30 ppm and 100 ppm fluoride (from NaF), respectively, in their drinking water during the last (3^{rd}) week of pregnancy and throughout the lactation period. The litters were separated from the mother rats on weaning and were then exposed to the respective levels of fluoride (0.5 ppm, 30 ppm, and 100 ppm) in drinking water up to the age of 10 weeks. The ten-week-old rats of control (n = 15), 30F (n = 13), and 100F (n = 9) groups were sacrificed after light ether anaesthesia. They were perfused transcardially with 0.9% saline followed by 10% formalin, and the brains were removed.

The brains were fixed in 10% formalin, and 5- μ m thick coronal sections of the hippocampus, amygdala, motor cortex, and cerebellum were taken. The sections were processed, and stained in 0.1% cresyl violet stain. The stained sections were observed under a binocular light microscope.

RESULTS AND DISCUSSION

No histological changes were seen in the brain regions of the 30F group when compared to the corresponding brain regions of the control group. The 100F group, however, showed significant changes in the histology of the various brain regions; the hippocampus, amygdala, motor cortex, and cerebellum showed neurodegenerative changes (Figures 1-6).

In the hippocampus, neurons were shrunken and darkly stained with a small nucleus, and there was a decrease in the cell number. Histological alterations were seen in cornu ammonis-1 (CA₁), cornu ammonis-4 (CA₄), and dentate gyrus sub regions of the hippocampus (Figures 1-3), but not in CA₂ and CA₃ sub regions. The hippocampus showed the most pronounced changes among the various brain regions of 100F rats. In rats exposed to long-term intake of high levels of fluoride in drinking water, Mullenix *et al*³ reported fluoride accumulation in important regions of the brain, especially the hippocampus. The pattern of changes seen in this study are similar to the findings of Varner *et al*⁴ who reported cellular abnormalities in the CA₁ and CA₄ areas of the hippocampus, and not in CA₂ and CA₃, in adult rats exposed to 1-ppm fluoride in drinking water for 52 weeks.

Neurodegenerative changes seen in the amygdala, motor cortex, and cerebellum were similar to those seen in the hippocampus (Figures 4-6). In the motor cortex, neurons in the superficial layers were the most affected, and there were also signs of chromatolysis and gliosis (Figure 5). Varner *et al*⁴ observed chromatin clumping, enhanced protein staining, pyknosis, vacuolation, presence of ghost-like cells, and decreased neuronal density in adult rats exposed to chronic intake of 1-ppm fluoride in drinking water. In our work, the Purkinje cells in the cerebellum were the most affected cell population, and there was an increase in the number of granular cells but a decrease in the number of Purkinje cells and molecular cells (Figure 6).

Fluoride 35 (1) 2002

pride (as g stages how any pm fluocampus, e in size Purkinje ne motor oride inentiation,

oxication,

rological remities, od-brain erimental ation influoride ar integain,⁵⁻⁷ in

orain cell oride in ology of

The rats dia), and r proper

nd 100F. the 30F

edical Colept. of Bioot. of Anat-Biochemis-



B

Figure 1. Photomicrographs of CA₁ region of hippocampus in control (A) and 100 ppm fluoride-treated (B). Note the degenerated neurons (arrows) in B. Cresyl violet stain, scale bar = $25 \ \mu m$.

Fluoride 35 (1) 2002

Figure 2. Photomic control (A) and 100 pp neurons (arrows) in



Figure 2. Photomicrographs of CA₄ region of hippocampus in control (A) and 100 ppm fluoride-treated (B). Note the degenerated neurons (arrows) in B. Cresyl violet stain, scale bar = 25 μm.

ated

Fluoride 35 (1) 2002

ė,





Figure 3. Photomicrographs of dentate gyrus region of hippocampus in control (A) and 100 ppm fluoride-treated (B). Note the degenerated neurons (arrows) in B. Cresyl violet stain, scale bar = 25 μm.







eaused marked train degeneral s may from the optimit heats it convicts patients, and distuiteres of the optimit of the optit of the optimit **Figure 6**. Photomicrographs of cerebellum in control (A) and 100 ppm fluoride-treated (B). Note the degenerated neurons (arrows) in B. Cresyl violet stain, scale bar = 25 µm.

Fluoride 35 (1) 2002

0

20 Shivarajashankara, Shivashankara, Bhat, Rao, Rao

Earlier, Zhavoronkov⁸ reported foci of demyelination in the cortex and subcortical areas, a decrease in the number of Purkinje cells in the cerebellum, swelling and irregular staining of the Nissl substance, and pyknosis of neurons in experimental animals subjected to fluoride intoxication.

In the present study, mother rats were exposed to high levels of fluoride in the last week of pregnancy and during lactation. This was done to ensure that the mother's body system became laden with fluoride so the litters would receive fluoride through the mother. Later, the litters were weaned and exposed directly to fluoride in drinking water. Thus, the young rats were subjected to chronic fluoride intoxication during the developing stages of life. Li *et al*⁹ reported a lower intelligence quotient (I.Q.) in children living in fluoride-endemic areas and suggested that the effect of exposure to high level of fluoride on intelligence probably occurs at an early stage of development of the embryo and infant when differentiation of brain cells is occurring and development is most rapid. Earlier, Mullenix *et al*³ observed abnormal behaviour patterns in rats exposed to intake of high levels of fluoride during prenatal, weanling, or adult stages. However, we did not investigate changes in the rat behaviour patterns that might have occurred in our study.

In conclusion, we find that chronic fluoride intoxication in the early stages of life caused marked neurodegenerative changes in the brain of rats. These changes may form the neural basis for impaired learning and memory, abnormal behaviour patterns, and disturbed overall body physiology.

ACKNOWLEDGEMENT

We are grateful to Dr B Mallikarjun, Dean, M R Medical College, Gulbarga, for his support and encouragement of this research project.

REFERENCES

- 1 Waldbott GL, Burgstahler AW, McKinney HL. Fluoridation: The great dilemma. Lawrence, Kansas: Coronado Press;1978.
- 2 Geeraerts F, Gijs G, Finne E, Crokaert R. Kinetics of fluoride penetration in liver and brain. Fluoride 1986;19:108-12.
- 3 Mullenix PJ, Denbesten PK, Schunior A, Kernan WJ. Neurotoxicity of sodium fluoride in rats. Neurotoxicol Teratol 1995;17:169-77.
- 4 Varner JA, Jensen KF, Horwath W, Issacson RL. Chronic administration of aluminium or sodium fluoride to rats in drinking water: alterations in neuronal and cerebrovascular integrity. Brain Res 1998;784:284-98.
- 5 Shashi A. Studies on alterations in brain lipid metabolism following experimental fluorosis. Fluoride 1992;25:77-84.
- 6 Shashi A, Singh JP, Thapar SP. Effect of long-term administration of fluoride on levels of proteins, free amino acids and RNA in rabbit brain. Fluoride 1994;27:155-9.
- 7 Vani ML, Reddy KP. Effect of fluoride accumulation on some enzymes of brain and gastrocnemius muscle of mice. Fluoride 2000;33:17-26.

Published by the Inte Editorial Office: 727 Brighton I

Fluoride 35 (1) 2002

- 8 Zhavoronkov AA. Non-ski 91.
- 9 Li XS, Zhi JL, Cao RO. Eff Fluoride 1995;28:189-92.

Histol

Histological changes in the brain in fluoride intoxication 21

cortex and he cerebelwknosis of

fluoride in to ensure the litters re weaned g rats were s stages of dren living ire to high e of devels is occurserved abof fluoride investigate our study.

arly stages rats. These emory, ab-

lege, Gul-

e great dinetration in

of sodium

istration of in neuronal

ing experi-

of fluoride n. Fluoride

nzymes of

Zhavoronkov AA. Non-skeletal forms of fluorosis. Arch Pathol 1977;39:83-91.

9 Li XS, Zhi JL, Cao RO. Effect of fluoride exposure on intelligence in children. Fluoride 1995;28:189-92.

30.0000 ARY: A pareiva dosimator realigned to persuits anappuredministration of the fluoride was leaved under laboration, conditions over the concentration was chosen in the closen of 0.16 – 4.47 mg FF/m². The forwelt concentration was chosen to be closen the Polish TLV (0.4 mg Hilm²O Timi dosimeter) burget of a Wistman N filter paper improgramed with K, MPO, placed in a polyethylene helder placed by a polyethylene screek Andre of the polyethylene helder of a state of a was chosen to be closen to closen to closen to closen to be closen to

reacting ecomposings are major environmental politicants, in remained rank seventh on the list of most dangerous chemical substances. In amounts of them are present in the work-place and in the groups of minima and magnesium foundries, phosphoric acid and phosphere twitt plums, petrolerum refinence, portery king, and factories where the ecomcusing, or welding is performed. The post common fluerus compopresent in the industrial davironment is hydrogen fluorde.

In order to prevent excessive expositing of workers to basic compocontinuous [nonitoring of the operpational environment is previous due the refineds used in monitoring of industrial polluting is previous does for obtainabilitie hable information on individual exposure to articreto taminants). This technique is used successfully to evaluate exceptant to the gen datas, suffit duotide, formatischyde, and some organic celeting, for duton to reports of active dosimeters for monitoring exposure to informafunction, ¹⁴ a passive device containing an alkali-impregnated polypropocollection element-insteameters for monitoring exposure to be the collection element-insteameters for monitoring exposure to be the second the second and an alkali-impregnated polypropo-

Passive dosinaters are appropriately selected and prepared with ents located in open holders attached to work clothane in the human The active selection on a volatile to the thetaintes firm the atmosphere selection of the selection of the distinctes may attach to the selection of the selection desimater, in which a selection device selection of the selection attaches and the device selection of the selection attaches and the device selection of the selection of the selection attaches and the selection of the selection of the device selection of the selection of the selection of the device selection of the selection of the selection of the are an even of the selection of the sel

Published by the International Society for Fluoride Research Editorial Office: 727 Brighton Road, Ocean View, Dunedin 9051, New Zealand