

EFFECTS OF FLUORIDE ION TOXICITY ON ANIMALS, PLANTS, AND SOIL HEALTH: A REVIEW

Surraya Shahab,^a Ghulam Mustafa,^{a,*} Imran Khan,^a Muhammad Zahid,^b Maimoona Yasinzai,^a
Nosheen Ameer,^a Nazia Asghar,^a Ikram Ullah,^a Akhtar Nadhman,^a Afaq Ahmed,^c
Iqbal Munir,^c Adnan Mujahid,^d Tajamal Hussain,^d Muhammad Nauman Ahmad,^e
Shaikh Saeed Ahmad^f

Islamabad, Pakistan

ABSTRACT: Substantial multi-disciplinary efforts have been made to investigate the effects of environmental fluoride ion (F) pollution since the last century. The chronic ingestion of high doses of F may adversely affect human health by causing skeletal fluorosis, dental fluorosis, bone fractures, the formation of kidney stones, decreased birth rates, weakening of thyroid functionality, and impair intelligence, particularly in children. High concentrations of F in soil may seriously threaten the life of plants, devastate soil microbial activity, disrupt the soil ecology, and cause soil and water pollution. In this review, we discuss the contribution of F to soil pollution and present certain remedies.

Key Words: Accumulation; Contamination; Dental fluorosis; Environment; Fluorosis; Pollutants; Remediation; Skeletal effects; Toxicity.

INTRODUCTION

The quality of life and the health of the environment are directly related to each other. The increasing incidence of animal and human health problems due to industrial pollution and anthropogenic environmental changes has attracted worldwide attention and efforts to find new remedies to better manage and sustain the environmental component.^{1,2} Toxic pollutants may be released to the environment via the air, soil, and water. Stack emissions to the air may add pollutants to the soil which may accumulate in plants via their roots. These pollutants accumulate in the food chain and then affect humans and wild life. Due to variations in natural and anthropogenic activities, global pollution is increasing and leading to contamination of the ecosystem with metals, non-metals, organic compounds, and inorganic compounds. The major contributors to this contamination are pesticides, sewage disposal, insecticides, herbicides, and the uncontrolled discharge of wastes. In the industrialized world, a large part of the population is exposed daily to a variety of chemicals and toxic metals which are harmful for human health. Compounds with the element fluorine are extensively utilized in almost every biological industry, and pollution by the fluoride ion (F) is widespread in the environment. Although F has an anticaries effect when applied topically to the teeth, fluorine is not an essential trace element and is not necessary for the development of healthy teeth and bones.^{3,4} Excessive F intake may adversely effect the health of humans, animals, and plants.⁵

^aSulaiman Bin Abdullah Aba-Al-Khail Center for Interdisciplinary Research in Basic Sciences, International Islamic University, Islamabad, Pakistan; ^bDepartment of Chemistry, Agriculture University, Faisalabad, Pakistan; ^cInstitute for Biotechnology and Genetic Engineering, The University of Agriculture, Peshawar, Pakistan; ^dDepartment of Chemistry, University of Punjab, Lahore, Pakistan; ^eDepartment of Agricultural Chemistry, The University of Agriculture, Peshawar, Pakistan; ^fFatima Jinnah Women University, Rawalpindi, Pakistan *For correspondence: Ghulam Mustafa, Sulaiman Bin Abdullah Aba-Al-Khail Center for Interdisciplinary Research in Basic Sciences, International Islamic University, Islamabad, Pakistan; E-mail: gmustafa@iiu.edu.pk.

Sources of F: There are many different sources of F but the main sources are the weathering of rocks, industrial emissions, and atmospheric deposition.⁶ The most common sources of F are mineral and geochemical stores and a large proportion of the discharge of F into subsoil water takes place through the degradation of rocks containing fluorine.⁷ F is among the more abundant elements in earth crust and is present in various rocks with a range of approximately 100–1000 µg/mg, with 625 µg/mg being a typical value. High concentrations of F are present in granites, felsic, quartz monzonites, syenites, biotite, and granodiorites. F-containing rocks such as muscovite, pegmatites, amphibolites, and biotite micas supply F to groundwater and soil by different processes such as soil forming and weathering.⁸ There are also anthropogenic sources of F such as the emission of hydrogen fluoride (HF) to the air or the addition of fluoride to water with various human activities, e.g., motorization, fluoridation of drinking water supplies, industrialization, and utilization of F-containing pesticides.⁹ Production of phosphate fertilizers is a major industrial source of F. A substantial amount of fluorine, e.g., 3.5%, is present in fertilizer made from rock phosphate but this percentage is reduced in the manufacturing process.⁸ Other anthropogenic sources for the entry of F into the earth include the current utilization of chemicals, such as phosphate manures, calcium fluoride (CaF), sodium fluoride (NaF), hydrogen fluoride, sulfur hexafluoride (SF₆), etc. When F is released into the air, it is carried by the wind to the surrounding soil and vegetation and contaminates them. Contamination of soil with F is basically due to utilization of F-containing fertilizers such as phosphorous fertilizers. The F content of soil commonly varies from 150–400 µg/mg and the value may increase in heavy clay soil up to 1000 mg/kg. The contaminated soil ultimately affects human beings through direct contact, the inhalation of vaporized soil, and the use of water contaminated with F by its passage through the soil. The use and production of phosphate fertilizers, the ceramic, zinc and steel industries, and energy plants are noteworthy sources of F pollution in the environment.¹⁰

Spatial distribution of F: The total concentration of F in the soil is usually derived from the parent material and therefore its distribution pattern in soil is related to the process of soil formation. The average content of F in the soil in world-wide has been estimated as 329 µg/mg. The lowest F content are usually present in sandy soil in relatively humid environments, while the higher concentrations of F are found in soil from weathered mafic rocks and in heavy clay soil. The pH of the soil, clay, and organic carbon content are the prime determinants of soil F content.¹¹ F enters the soil through different ways such as dry deposition, precipitation, and with contaminated litter where it is absorbed readily. The absorbed F increases the total soluble F concentration in the soil, influences the pH of soil, and can combine with toxic elements such as aluminum and heavy metals. F can exist as the free fluoride ion (F⁻) or form complexes with elements such as iron (Fe), boron (B), calcium (Ca), and aluminium (Al), with the complexes of Al and F being most prevalent.¹²

Effects of F pollution: The aerial F emitted from industries not only pollutes the air but also, along with ground water contaminated by fluorine-containing mineral deposits, contaminates plants, crops, soil, vegetables, and freshwater bodies. The use

these F-contaminated products and sources may result in toxic effects on health of both humans and domestic animals.¹³ While the topical use of F on teeth is regarded as having beneficial effects, the evidence for any beneficial effects from systemic absorption is now considered to be weak.⁴

Impact of F on the human system: The research has shown repeatedly that the consumption of F can extremely be harmful and in some cases deadly. In terms of acute toxicity, F is slightly less toxic than arsenic and more toxic than lead. High concentrations of F can lead to serious poisoning incidents with death.




Figure 1. Metabolism of F in the human body.¹³⁻¹⁶

Chronic toxicity can occur with long term exposure. Absorption from the gastrointestinal tract and lungs occurs rapidly with the peak in the plasma occurring after 30 min. Fluoride is cleared from plasma through uptake by bones and excretion in the urine with infants and children clearing relatively larger amounts of F into bones because of their growing skeleton.¹⁴ After a single dose, plasma concentrations rise to a peak and then fall as the F is cleared by the renal system and bone back to a short-term baseline with a half-life of several hours.¹⁴ For each mg of F that is ingested approximately 0.9 mg is absorbed in the stomach and small intestine and passes to the blood while 0.1 mg is excreted unabsorbed in faeces, 0.40 mg is stored in bones and teeth, 0.45 mg is excreted in the urine, and 0.05 mg is excreted in breast milk, sweat, and saliva.^{14,15} In the rat, approximately 25% of F absorption occurred in the stomach and 75% from the small intestine.¹⁶ The skeleton and teeth are the prime organs of F retention/accumulation in human body, while relatively small amounts may be deposited in another calcifying organ, the pineal gland.^{14,17} The retention period of F can be up to several years. F can reach the foetus via the placenta.⁹ (Figure 1).

Fluorosis: A high consumption of F may cause chronic fluorine toxicity, known as fluorosis¹. One of the mechanisms involved in the pathogenesis of fluorosis is increased oxygen radical generation and lipid peroxidation. Dental fluorosis, with

mottling of the teeth, is common in F endemic areas. The teeth develop chalky white patches and become rough. In addition, yellow to dark brown lines may become visible.¹⁷ Yellow to brown, or even become black, opaque patches may occur with more severe toxicity with increased tooth porosity which leads to chipping or pitting.¹⁸ In skeletal fluorosis the mass and density of bone increases along with the development of skeletal and joint symptoms. Severe pain in the spine, joints, and hip area are early symptoms. The symptoms of “poker back” may appear with increasing stiffness until whole spine converts into one continuous column of bone. A number of muscles of the spine may also become calcified and paralyzed as the condition increases in severity. Neurological defects, complete paralysis, crippling deformities of the spine and the major joints, muscle wasting, and increased density of the spinal cord may occur in the most advanced stages of skeletal fluorosis. Skeletal fluorosis may occur with a high F intake from water and other dietary factors.¹⁹




Figure 2: The transportation of fluoride from agricultural soil and irrigation water to plants, the consumption of grains containing fluoride by humans, and the accumulation of fluoride in humans with the development of dental and skeletal fluorosis.¹

Phytotoxicity of F: Fluorine exists in the environment as gaseous molecules (F_2) and in its reduced form as the ion fluoride (F). Under certain circumstances of comparatively lower pH and hardness fluorine occurs as the fluoride ion in water. Plants are exposed to F through different sources such as air, water, and soil. Natural sources of F include cryolite, apatitite, feldspar, volcanic gases, and marine aerosols.

Other important sources of F include the production of bricks, glass, aluminium, ceramics, and high phosphate fertilizers. The trace amounts of F available to plants by diffusion in the soil may be absorbed by roots. In plants such as tea, F is naturally accumulated. In some cases, F unfavorably affects the plant leaves after it intrudes into the leaves because of its high solubility. F deposition may become relatively slower over time. Fluoride may stop photosynthesis and other essential processes in plants. F passes from roots to leaves through the process of transpiration or moves through stomata and accumulates in the margins of the leaves. Marginal and tip necrosis is the first symptom of fluorine injury in plant leaves. However, such symptoms may also appear in drought or salinity stress, which can resemble injury caused by F. Unfortunately, an enormous number of plants are highly F sensitive such as dracaena, Tahitian Bridal Veil, the spider plant, etc. Economically important fruits are also sensitive to F toxicity such as apricot, peaches, plums, etc. Flowers may also be affected by F toxicity, e.g., tulip, gladiolus, lily, etc.²⁰ Arora and Bhateja reported that F in soil, crops such as wheat, rice, and potato, and in other dietary sources can lead to the occurrence of fluorosis. Therefore, one should carefully calculate the amount of F in daily diet.²¹




Figure 3. Mean F concentrations (ppm) of different crops in relation to the mean F concentration in the soil.²¹

Mishra et al. reported that accumulation of F in plant tissues is directly proportional to the amount present in soil.²² A number of these plants were examined and found to have a substantially decreased Net Primary Productivity (NPP) [above ground biomass (AGP) and below ground biomass (BGP)] and yield (pod weight) with exposure to F (Figure 4).

When treated with F in a concentration of 20–100 ppm, the NPP decreased in brinjal, mung, and tomato plants by 6.64–56.72%, 10.27–53.61%, and 14.46–62.24%, respectively. When treated with F in a concentration of 10–50 ppm, mustard, okra, and chillies also showed considerable declines in NPP of 15.58–61.21%, 12.28–52.78%, and 40.8–90.65%, respectively. When treated with F in a concentration of 20–100 ppm, the monocots maize and rice showed a decline in the NPP of 12.17–61.20% and 6.64–56.72%, respectively. Hence, it has been found that F toxicity in soil may harshly effect the NPP in several crops in minute as well as in extreme concentrations.²²

Reduction, compared to controls, in net primary production (NPP) and yield (%)




Figure 4. Percent reduction, compared to controls treated with tap water, in net primary production (NPP) [above ground biomass (AGP) and below ground biomass (BGP)] and yield (pod weight) in different crops treated with various concentrations of fluoride ion (F). A 100% reduction in yield indicates that no pods were formed.²²

- 1 = brinjal (F concentrations used in the F-treated groups: NPP=100 ppm, yield=100 ppm)
- 2 = tomato (F concentrations used in the F-treated groups: NPP=100 ppm, yield=1000 ppm)
- 3 = mustard (F concentrations used in the F-treated groups: NPP=50 ppm, yield=50 ppm)
- 4 = mung (F concentrations used in the F-treated groups: NPP=100 ppm, yield=100 ppm)
- 5 = ladies finger (F concentrations used in the F-treated groups: NPP=50 ppm, yield=50 ppm)
- 6 = maize (F concentrations used in the F-treated groups: NPP=100 ppm, yield=20 ppm)
- 7 = paddy (F concentrations used in the F-treated groups: NPP=100 ppm, yield=100 ppm)
- 8 = chilli (F concentrations used in the F-treated groups: NPP=100 ppm, yield=20 ppm)

Due to the high capacity of some plants for the uptake of F from soil the monitoring of soil F levels is necessary from time to time. Spinach is a popular vegetable and has a high capacity for taking up F intake so that this plant may have high F levels and consequently there may be an increased risk of health consequences after its consumption. Therefore, it is required that this type of vegetable be cultivated away from soil that has accumulated high F levels.²³ The accumulation and uptake of F in the shoot and root of spinach (*Spinacea oleracea*) and the effects of contamination of soil by inorganic F (NaF) were investigated in a pot experiment under controlled conditions.²³ Using soil with a soluble F range of 2.57–16.44 mg/kg soil, the experimental results showed that most of the F accumulated in the spinach tissues and the plant had a mechanism for partitioning the water labile F and the total F in its tissues (Figure 5). It restricted the translocation of F from the root to the shoot, thus sparing the part of the plant mostly used for consumption.²³




Figure 5. Effect of NaF on the F uptake of *Spinacea oleracea*. LSD (0.05): Least significant difference between the means at the 0.05 probability level: shoot 54.7, root 21.3.²³

Several plants have the capability to accumulate soil F through several mechanisms. The potato plant has a high capacity to accumulate soil F and the strong fluorine accumulation in the shoots and leaves indicates the potential of *S. tuberosum* for field application for the removal of fluorine. In the case of the potato, the tolerance index and growth ratio have different trends from each other as concentration of F changes. The F accumulation is highest in the leaves while percentage total F translocation from soil to plant linearly decreases as the soil F increases. Therefore, the potato (*S. tuberosum*) can be a suitable candidate species for the removal of F for phytoremediation purposes.²⁴




Figure 6. Fluoride accumulation (mg F/kg potato) in potato roots, shoots, leaves, and bulbs at different levels of soil fluoride. The bars represent the least significant difference (LSD) between the means at the 0.05 probability difference.²⁴

Tea plants have a high ability to absorb F from soil and surrounding air and accumulate in the leaves in the form of an Al and F complex. Consumption of high F tea for long time can result in chronic F intoxication.²⁵

Toxicity of F in birds and animals: The fluoride ion (F) is not considered to be essential for human growth and development,⁴ including for the development of healthy teeth and bones, and the chronic ingestion of fluoride at high levels (above 6 mg/day) can be toxic to animals and human beings and cause dental, skeletal, and non-skeletal fluorosis.²⁶ Fluorosis usually occurs in two forms: (i) endemic fluorosis caused by drinking water or consuming food with a high F content and (ii) industrial fluorosis resulting from exposure to air containing a considerable F content. Fluorosis, in humans, animals, and birds, affects not only the skeletal parts of the body but also affects soft tissues, e.g., brain, liver, kidney, thyroid, and spinal cord. Anjum et al. investigated the effect of a high concentration of F on hepatic and renal enzymes in four groups of domestic chickens, A, B, C, and D, receiving 0, 10, 20, and 30 µg/g of NaF by body weight, respectively, on a weekly basis for four weeks.²⁷ Alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT), and bilirubin were determined as indicators of liver function, while uric acid was used as a parameter for renal function. The results showed high values for all the parameters ($p < 0.05$) in the F-treated groups indicating that F effected hepatic and renal function in the exposed birds.²⁷

Although all animals are susceptible to high doses of F, the tolerance level changes from one species to another. Noteworthy sources of F for terrestrial animals are drinking water, soil, and vegetation contaminated with F emitted by different activities such as volcanic eruptions and industrial activities²⁸ (Figure 7). The metabolism of F in animals is similar to that of humans²⁸ (Figure 8).

Among the terrestrial vertebrates, herbivores are more susceptible than carnivores and other animals. Domestic and wild herbivores are more exposed to environmental F contamination because they are nonselective eaters and can consume contaminated feed, water, and forage. Cattle and sheep have attracted more attention from researchers worldwide, perhaps due to their large populations and their greater economic importance. However, other animals, including water buffaloes, horses, goats, pigs, and wild cervids, can also suffer from F-toxicity naturally.²⁸ Skeletal, non-skeletal, and dental fluorosis have been studied in buffaloes (*Bubalus bubalis*), camels (*Camelus dromedarius*), donkeys (*Equus asinus*), horses (*Equus caballus*), and cattle (*Bos taurus*).²⁹

Choubisa et al. studied F toxicity in 760 domestic animals comprising 386 cattle (*Bos taurus*), 158 goats (*Capra hirus*), 7 donkeys (*Equus asinus*), 11 horses (*Equus caballus*), 131 buffaloes (*Bubalus bubalis*), and 67 sheep (*Ovis aries*) from different villages with a mean water F range of 1.5–4.4 ppm.³⁰ Three hundred and seventy (48.7%), were found to be afflicted with dental mottling. On clinical examination, 325 (42.8%) of the animals revealed periosteal exostoses, intermittent lameness, hoof deformities, stiffness in the legs and tendons, and wasting of the main mass of the hind quarter and shoulder muscles. Fluorosed animals also showed signs of non-skeletal fluorosis such as gastro-intestinal discomforts, impaired reproductive

functions, neurological disorders, and congenital abnormalities. The prevalence and severity of these F effects increased with an increasing F concentration in the water




Figure 7. Sources of fluoride in animals.²⁸




Figure 8. Fluoride metabolism in the animal body after its oral ingestion.²⁸

Remediation of F-contaminated soils: Moon et al. described a method for the removal of F from contaminated soil by washing with different solutions such tartaric acid ($C_4H_6O_6$), nitric acid (HNO_3), sulfuric acid (H_2SO_4), sodium hydroxide ($NaOH$), and hydrochloric acid (HCl) (Figures 9–11).³¹ The concentrations of the washing solutions ranged from 0.1 to 3 M with a liquid to solid ratio of 10. The results of washing with the various solutions indicated that HCl was most the effective for removing F from contaminated soils. The highest F removal rate from the contaminated soil, of approximately 97%, was obtained using 3M HCl. The F-

removal efficiencies of the washing solutions were in the following order:
 $\text{HCl} > \text{HNO}_3 > \text{H}_2\text{SO}_4 > \text{NaOH} > \text{C}_4\text{H}_6\text{O}_6$.³¹




Figure 9. Fluoride concentrations remaining in the soil after hydrochloric acid (HCl) washing at various concentrations together with the Korean warning standard for one area.³¹




Figure 10. Fluoride concentrations remaining in the soil after nitric acid (HNO_3) washing at various concentrations together with the Korean warning standard for one area.³¹




Figure 11. Fluoride concentrations remaining in the soil after tartaric acid ($C_4H_6O_6$) washing at various concentrations together with the Korean warning standard for one area.³¹






Figure 12. Fluoride concentrations remaining in the soil after sulfuric acid (H_2SO_4) washing at various concentrations together with the Korean warning standard for one area.³¹

Zhu et al. investigated the electrokinetic remediation of F-contaminated field soil with an organic content, pH, and initial F-concentration of 20.52 g/kg, 18.17, and 1058 mg/kg, respectively.³² Electrokinetic experiments were carried out under two different concentrations of alkaline solution and three different voltage gradients (Figures 12A-12D). The removal efficiency of fluorine increased up to 73% within 10 days with increasing the concentration of the alkaline solution and the applied voltage. This process could effectively promote the migration of F present in soil. The main transport mechanism was electromigration and the electroosmotic flow had an effect on the soil F migration. An appropriate anolyte enhanced electrokinetic method can be applied to remove fluorine from contaminated field soil and also has a significant potential for removing other anionic pollutants such as arsenic and chromium from soil.³²




12A




12B

Figures 12A and 12B. 12A: Schematic diagram of the electrokinetic apparatus; 12B: Soil electrical conductivity (EC, $\mu\text{s}/\text{cm}$) in soil sections after the electrokinetic treatments.³²




12C




12D

Figures 12C and 12D. 12C: Cumulative mass of fluorine in the anolyte (mg); 12D: Cumulative mass of fluorine in the catholyte (mg).³²

Zhou et al. reported the effect of changing the pulse interval of an electric field on the remediation of fluorine-contaminated soil by an electrokinetic remediation method.³³ The experiment results indicated that at the same intensity of the electric field, pulse-enhanced electrokinetic remediation showed a better performance in the removal of fluorine than conventional electrokinetic remediation. The efficiency of the pulsed enhanced electrokinetic remediation was increased because concentration polarization decreased with applying the electric field and it increased the electric current, the electrical voltage, and the electroosmotic flow in the soil cell. Therefore, for the removal of fluorine from soils, a pulse-enhanced technique would be a preferred choice.³³




13A



13B

Figures 13A and 13B. 13A: A schematic diagram of the experimental apparatus. Key to numbering: 1. The DC power supply (GPC6030D, Gw instek, China); 2. The soil cell (10 [L] cmx6 [W] cmx8 [H] cm); 3, 4. The electrode compartment (4 [L] cmx6 [W] cmx8 [H] cm); 5, 6. Working electrode (graphite sheet, 1 [L] cmx6 [W] cmx8 [H] cm); 7. Anion exchange membrane (3361BW, Shanghai Shinghua Water Treatment Material Co., Ltd.); 8. Cation exchange membrane (3362BW, Shanghai Shinghua Water Treatment Material Co., Ltd.); 9, 10. The electrolyte reservoir; 11-14. 4-Channel peristaltic pump (BT00-300T/DG-4, longerpump, China); 15, 16. Simple flow regulator; 17. Time; 13B: Residual fluorine concentration in soil sections after electrokinetic remediation.³³

Kim et al. studied electrokinetic remediation for the removal of fluorine from contaminated soil and the results showed that electro kinetic techniques increased the efficiency of fluorine removal by up to 75.6%.³⁴ From these results it can be concluded that analyte conditioning is a very effective enhancement method to remove fluorine from contaminated soil in electrokinetic remediation. This method can be also applied for removal of anionic pollutants such as chromate and arsenic from contaminated soil.



Figures 14A and 14B. Schematic diagram and dimensions of the experimental apparatus: (A) anolyte circulation system; (B) dimensions of the experimental apparatus.³⁴

CONCLUSIONS

This review considers the importance of F and argues that a high intake of F, via ingestion or inhalation from a variety of sources, may cause toxicity in humans and animals, including dental, skeletal, and non-skeletal fluorosis. The F toxicity can be acute or chronic depending on the level and the duration of the F-intake. The studies in the literature show that the presence of a high concentration of F in soil affects plants and aquatic life and leads to soil and water pollution. Plants species with a susceptibility to F pollution in soil may be drastically damaged. In addition, F pollution may have a devastating effect on the microbial activity in soil and disrupt the soil ecology.

REFERENCES

- 1 Ahmed DE. Environmental pollution and its impact on human and animal health: a review The Sudan J Vet Res 2007;22:37-46.
- 2 Tsiros TX, Haidouti C, Chronopoulou A. Airborne F contamination of soil and olive trees near an aluminum plant: measurement and simulation. J Environ Sci Health 1998;33(7):1309-24.
- 3 Chouhan S, Flora SJ. Arsenic and F: two major ground water pollutants. Indian Journal of Experimental Biology 2010;48:666-78.
- 4 Scientific Committee on Health and Environmental Risks (SCHER). Opinion of critical review of any new evidence on the hazard profile, health effects, and human exposure to fluoride and the fluoridating agents of drinking water. Brussels, Belgium: Directorate General for Health and Consumers, European Commission; 2011 May 16. pp. 2-4.
- 5 Koblar A, Tavcar G, Ponikvar-Svet M. Effects of airborne F on soil and vegetation. Journal of Fluorine Chemistry 2011;132:755-75.
- 6 Xiaojing L, Binbin W, Baoshan Z. Geochemical process of fluorine in soil. Chin J Geochem 2014;33:277-9.
- 7 Jacks G, Bhattacharya P, Chaudhary V, Singh PK. Controls on the genesis of some high-F groundwaters in India. Applied Geochemistry 2005;20:221-8.
- 8 Bhat N, Jain S, Asawa K, Tak M, Shinde K, Singh A, et al. Assessment of fluoride concentration of soil and vegetables in vicinity of zinc smelter, Debari, Udaipur, Rajasthan. J Clin Diagn Res 2015;9(10):ZC63-6.
- 9 Ghosh A, Mukerjee KS, Saha B. Sources and toxicity of fluoride in the environment. Res Chem Intermed 2013;39:2881-915.
- 10 Paul ED, Gimba CE, Kagbu JA, Ndukwe GI, Okibe FG. Spectrometric determination of fluoride in water, soil and vegetables from the precinct of river Basawa, Zaria, Nigeria. J Basic Appl Chem 2011;1:33-8.
- 11 Kumar B, Naaz A, Shukla K, Narayan C, Singh G, Kumar A, et al. Spatial variability of fluorine in agricultural soils around Sidhi District, Central India. Journal of Geological Society of India 2016;87(2):227-35.
- 12 Domingos M, Klumpp A, Rinaldi MCS, Modesto IF, Klumpp G, Delitti WBC. Combined effects of air and soil pollution by fluoride emissions from *Tibouchina pulchra* Cogn., at Cubatão, SE Brazil, and their relations with aluminium. Plant and Soil 2003;249:297-308.
- 13 Choubisa SL, Choubisa D. Status of industrial fluoride pollution and its diverse adverse health effects in man and domestic animals in India. Environ Sci Pollut Res 2016;23:7244-7254.
- 14 Doull J, Boekelheide K, Farishian BG, Isaacson RL, Klotz JB, Kumar JV, Limeback H, Poole C, Puzas JE, Reed N-MR, Thiessen KM, Webster TF, Committee on Fluoride in Drinking Water, Board on Environmental Studies and Toxicology, Division on Earth and Life Studies, National Research Council of the National Academies. Fluoride in drinking water: a scientific review of EPA's standards. Washington, DC: The National Academies Press; 2006. p.181-204. [Contract No.: 68-C-03-013. Sponsored by the U.S. Environmental Protection Agency]. Available to read or purchase on line at www.nap.edu. pp. 91, 92, 100, and 253-6.
- 15 Limeback H, Robinson C. Fluoride therapy. In: Limeback H, editor. Comprehensive preventive dentistry. Ames, Iowa, USA, and Chichester, West Sussex, UK: Wiley-Blackwell, an imprint of John Wiley & Sons; 2012. pp. 251-82.
- 16 Nopakun J, Messer HH, Voller V. Fluoride absorption from the gastrointestinal tract of rats. J Nutr 1989;119(10):1411-7.
- 17 Kumar N, Bansal N, Sharma SK. Determination of fluoride status in ground water of Rajasthan. IJPCBS 2014;4:576-92.
- 18 David L, Ozsvath. Fluoride and environmental health: a review. Rev Environ Sci Biotechnol 2009;8:59-79.
- 19 Annaduraia TS, Rengasamy KJ, Sundaramb R, Munusamy PA. Incidence and effects of fluoride in Indian natural ecosystem: a review. Advances in Applied Science Research 2014;5:173-85.
- 20 Kumar B, Anshumali. Fluoride in agricultural soil: a review on its sources and its toxicity to plants. Global sustainability transitions: impacts and innovations. ISBN 978-93-83083-77-0. pp. 29-37. Available from: http://www.krishisanaskriti.org/vol_image/10Sep201505094305.pdf
- 21 Arora G, Bhateja S. Estimating the fluoride concentration in soil and crops grown over it in and around Mathura, Uttar Pradesh, India. American Journal of Ethnomedicine 2014;1:036-041.
- 22 Mishra CP, Sahu KS, Bhoi KA, Mohapatra CS. Fluoride uptake and net primary productivity of selected Crops. Open Journal of Soil Science 2014; 4: 388-398
- 23 Jha SK, Nayak KA, Sharma KY. Response of spinach (*Spinacea oleracea*) to the added F in an alkaline soil. Food and Chemical Toxicology 2008;46:2968-71.
- 24 Das C, Dey U, Chakraborty D, Datta JK, Mondal NK. Fluoride toxicity effects in potato plant (*Solanum tuberosum* L.) grown in contaminated soils. Octa Journal of Environmental Research 2015;3(2):136-43.
- 25 Yi J, Cao J. Tea and fluorosis. Journal of Fluorine Chemistry 2008;129:76-81.
- 26 World Health Organization. Fluorides: environmental health criteria 227. Geneva: World Health Organization (Published under the joint sponsorship of the United Nations Environment Programme, the International Labour Organization and the World Health Organization, and produced within the framework of the Inter-Organization Programme for the Sound Management of Chemicals); 2002. Available from: <http://www.inchem.org/documents/ehc/ehc/ehc227.htm#1.5>
- 27 Anjum MK, Mughal SM, Sayyed U, Yaqub A, Khaliq A, Rashid AM, et al. Influence of increasing fluoride dose rates on selected liver and kidney enzymes profile in domestic chicken (*Gallus domesticus*). Journal of Animal & Plant Sciences 2014;24(1):77-80.
- 28 Ranjan R, Ranjan A. Fluoride toxicity in animals. Springer Briefs in Animal Sciences (part of an E-book collection). Basel, Switzerland: Springer International Publishing; 2015. pp. 7-12.
- 29 Choubisa SL. Natural amelioration of fluoride toxicity (fluorosis) in goats and sheep [scientific correspondence]. Current Science 2010; 99(10):1331-2.
- 30 Choubisa LS, Mishra VG, Sheikh Z, Bhardwaj B, Mali P, Jaroli VJ. Toxic effects of fluoride in domestic animals. Adv Pharmacol Toxicol 2011;12:29-37.
- 31 Moon DH, Jo R, Koutsospyros A, Cheong KH, Park JH. Soil washing of fluorine contaminated soil using various washing solutions. Bull Environ Contam Toxicol 2015;94:334-9.
- 32 Zhu S, Zhang, Dong T. Removal of fluorine from contaminated field soil by anolyte enhanced electrokinetic remediation. Environ Earth Sci 2009;59:379-84.
- 33 Zhou M, Zhu S, Liu F, Zhou D. Pulse-enhanced electrokinetic remediation of fluorine-contaminated soil. Korean J Chem Eng 2014; 31:2008-13.
- 34 Kim DH, Jeon CS, Baek K, Ko SH, Yang JS. Electrokinetic remediation of fluorine-contaminated soil: Conditioning of anolyte, Journal of Hazardous Materials 2009;161:565-9.